

TITLE

A pan-TE map highlights transposable elements underlying domestication and agronomic traits in Asian rice

AUTHORS

Xiaoxia Li^{1,2#}, Xiaofan Dai^{1,2#}, Huiying He^{1#}, Yang Lv^{1,2#}, Longbo Yang^{1#}, Wenchuang He¹, Congcong Liu^{1,3}, Hua Wei¹, Xiangpei Liu¹, Qiaoling Yuan¹, Xianmeng Wang¹, Tianyi Wang¹, Bintao Zhang¹, Hong Zhang¹, Wu Chen¹, Yue Leng¹, Xiaoman Yu^{1,4}, Hongge Qian¹, Bin Zhang^{1,3}, Mingliang Guo¹, Zhipeng Zhang¹, Chuanlin Shi¹, Qianqian Zhang¹, Yan Cui¹, Qiang Xu¹, Xinglan Cao¹, Dandan Chen¹, Yongfeng Zhou^{1,5*}, Qian Qian^{1,3,4*}, Lianguang Shang^{1,3*}

ADDRESS

¹Shenzhen Branch, Guangdong Laboratory of Lingnan Modern Agriculture, Genome Analysis Laboratory of the Ministry of Agriculture and Rural Affairs, Agricultural Genomics Institute at Shenzhen, Chinese Academy of Agricultural Sciences, Shenzhen, 518120, China

²Rice Research Institute, Shenyang Agricultural University, Shenyang 110866, China

³Yazhouwan National Laboratory, No. 8 Huanjin Road, Yazhou District, Sanya City, Hainan Province, 572024, China

⁴State Key Laboratory of Rice Biology, China National Rice Research Institute, Hangzhou, 310006, China

⁵National Key Laboratory of Tropical Crop Breeding, Tropical Crops Genetic Resources Institute, Chinese Academy of Tropical Agricultural Sciences, Haikou, 571101, China

These authors contributed equally to this work.

*Correspondence: shanglianguang@caas.cn, qianqian188@hotmail.com, zhouyongfeng@caas.cn.

ABSTRACT

Transposable elements (TEs) are ubiquitous genomic components and hard to study due to their highly repetitive. Here we assembled 232 chromosome-level genomes based on long-reads sequencing data. Coupling with 15 existing assemblies, we developed a pan-TE map comprising

both cultivated and wild Asian rice. We detected 177,084 high-quality TE variations and inferred their derived state using outgroups. We found TEs were one source of phenotypic variation during rice domestication and differentiation. We identified 1,246 genes whose expression variation was associated with TEs but not SNPs, such as *OsRbohB*, and validated *OsRbohB*'s relative expression activity using dual-LUC reporter assays system. Our pan-TE map allowed us to detect multiple novel loci associated with agronomic traits. Collectively, our findings highlight the contributions of TEs to domestication, differentiation, and agronomic traits in rice, and there is a massive potential for gene cloning and molecular breeding by the high-quality Asian pan-TE map we generated.

Keywords: transposable element; super pan-genome; pan-TE; rice

INTRODUCTION

Transposable elements (TEs) are ubiquitous components of genomes and one of the major forces driving genome evolution [1]. As early as 1948, Barbara McClintock observed that transposons were associated with color variegation in maize kernels and leaves [2], revealing the crucial role of TEs in phenotypic variation. TEs also represent a major source of genomic variation. Two classes of TE are roughly distinguished: retroelements and DNA transposons. Retroelements are further classified as long terminal repeats (LTRs, including *Gypsy*, *Copia*, and unknown elements) and non-LTRs (short and long interspersed retrotransposable elements [SINE and LINE, respectively]), and DNA transposons are classified as miniature inverted repeat transposable elements (MITEs, including *Stowaway* and *Tourist*), DTC (*CACTA*), DTA (*hAT*), DTT (*Tc1-Mariner*), DTM (*Mutator*), DTH (*PIF-Harbinger*), and *Helitron* elements [3]. Increasing evidence has supported the hypothesis that TEs are responsible for many well-known human diseases [4] and various plant phenotypes [5-7]. For example, a LTR retrotransposon insertion upstream of an apple MYB transcription factor, *MdMYB1*, a core transcriptional activator of anthocyanin biosynthesis, is associated with red-skinned phenotype of apple [8]. An insertion of a *Gret1* LTR retrotransposon in the Chardonnay grape variety, which is not present in the Cabernet variety, generated a loss-of-function allele of the *Vvmyb1A* gene, leading to a loss of color in the fruit [9]. A TE inserted in a regulatory region of the maize domestication gene (*tb1*) acts as an enhancer of gene expression and partially explains the increased apical dominance in maize compared to its

progenitor, teosinte [10].

Recently reported crop TE variation datasets mainly consist of TE variations obtained from Illumina short-read sequencing [5, 7, 11], such as one constructed using short reads sequences from 524 *Brassica rapa* accessions [5] and one constructed using short reads sequences from 602 tomato accessions [7]. However, identifying highly repetitive sequences such as TEs using short-read sequencing data is difficult and predictions have low accuracy, especially for some TE families that have small-length and a very high copy number, such as the SINE, LINE, and MITE families. Identification of TE variations using high-quality genome sequences based on long-read sequencing data is the most accurate method to identify TE variations. At present, a population-scale TE variation dataset based on de novo long-read assemblies has been published only in *Drosophila nasuta* [12].

Rice is a major food crop for half of the global population [13, 14], and improving rice productivity is essential to meet the growing demands of the ever-increasing world population [15, 16]. Asian cultivated rice (*Oryza sativa*, *Os*) was domesticated from Asian wild rice (*O. rufipogon*, *Or*) and has two main types: Geng (*Os. japonica*, *Osj*) and Xian (*Os. indica*, *Osi*) [17]. TEs are known to contribute to phenotypic variation in rice populations. For instance, a *Helitron* in the *MYB61* [18] promoter was reported to affect rice nitrogen utilization; an LTR insertion in the *GY3* promoter suppresses its expression and results in a higher content of active cytokinins in young panicles [19]; and the *PIF-Harbinger* transposon-derived gene *PANDA* epigenetically coordinates panicle number and grain size in rice [20]. Identification of TEs across rice accessions would facilitate efforts to determine the contribution of TEs to expression and phenotypic variation. Three studies of population-scale TE variations based on Illumina short-reads [11, 21, 22] have been published in rice. However, these studies mainly focused on *Os* accessions, with *Or* and domestication-associated TE variations remaining underexplored. In addition, although an increasing number of rice studies based on high-quality assemblies have been published, population-level, high-quality studies of TE variations have not been reported. Moreover, because of the absence of rice outgroup accessions in previous studies, it is hard to distinguish the ancestral and derived state of TE variations. Thus, building a pan-TE resource using high-quality chromosome-level assemblies and inferring whether TE variations are ancestral or derived based on outgroups are desirable, and such information will help us to know the true extent of TE

variation in plants and their biological impacts on rice environmental adaptation and domestication.

Here, we de novo assembled chromosome-level genomes for 232 diverse rice accessions by integrating Oxford Nanopore Technology (ONT) long-read data and Illumina short-read data from public databases. Combining these assemblies with 15 existing rice assemblies, we constructed a pan-TE map of 247 Asian accessions (218 *O.s* and 29 *O.r* accessions) and inferred whether TE variations were ancestral or derived using three non-Asian accessions as outgroups (*O. glaberrima*, *O. barthii*, and *O. glumaepatula*). We analyzed TE variation characteristics and the role of TE mobilization in rice domestication diversification. Moreover, we analyzed the contribution of TE variation to gene expression variation and identified 1,246 genes whose expression levels were significantly associated with TE variation but not single-nucleotide polymorphisms (SNPs). Importantly, our pan-TE map enabled genome-wide association study (GWAS)-based identification of loci associated with phenotypes (i.e., seed setting rate under cold stress), which were undetectable when using only SNPs. Taken together, the pan-TE map is a valuable resource for understanding the genetic basis of environmental adaptation, domestication, and agronomic traits in rice.

RESULTS

De novo assembly and pan-TE map construction for Asian rice accessions

We collected the relevant long-read and short-read data for 250 rice accessions, chromosomal-level assemblies for 18 accessions, and leaf transcriptome data for 202 *O.s* accessions, which were selected based on the geographical distribution and genetically diverse of rice from public databases (Supplementary **Table S1a**, See “**Materials and data collection**”). The 250 accessions included 247 Asian rice accessions and three non-Asian rice accessions (one *O. glaberrima*, one *O. barthii* and one *O. glumaepatula*) as outgroups (henceforth termed “outgroups”) (Supplementary **Table S1a**). A phylogenetic tree constructed using whole genome SNP data for the 250 accessions clearly divided these materials into four main subpopulations, *Osi*, *Aus*, *Osj*, and *Or* (**Fig. 1a**). Since data for only five *Aus* accessions were collected, which is not enough for population analysis, they were not included in subsequent subpopulation analysis. Finally, 146 *Osi*, 61 *Osj*, and 27 *Or* were retained to represent the *Osi*, *Osj*, and *Or* subpopulations for

population comparison analysis. The results of admixture and principal component analysis (PCA) were highly consistent with those of the phylogenetic tree (Supplementary **Fig. S1a-c**).

We found a genome has an average of 28,895 intact TEs (Supplementary **Fig. S1d**), and the number of intact TE of chromosomal-level genomes is more than that of contig-level genome (Supplementary **Fig. S1d**), thus obtaining high-quality chromosomal-level genomes was critical for constructing a pan-TE map. To this end, the non-chromosome-level assemblies of 232 accessions were de novo assembled with ONT long reads using NextDenovo (version 2.4) [23] (**Fig. 1b**), and the assemblies were corrected with both long reads and Illumina short reads using Nextpolish (version 1.4.0) [24]. The 232 new assemblies had an average length of 390.7 ± 11.3 Mb and average contig N50 of 23.7 ± 6.0 Mb (Supplementary **Table S1b**). An average of $99.7\% \pm 0.2\%$ of contigs were anchored to the chromosomes (Supplementary **Table S1b**). We next evaluated the quality of the new assemblies and found that: (i) an average of $98.5\% \pm 0.5\%$ of Illumina short reads could be mapped to its corresponding assembly, which is similar to the mapping rate (98.0%) obtained for Nipponbare (NIP) [15, 25] (Supplementary **Table 1b** and **Fig. S1e**); (ii) the assembly quality was comparable to that of the NIP genome [15, 25], with $97.9\% \pm 0.2\%$ of a Benchmarking Universal Single-Copy Orthologs (BUSCO) [26] 1,440 reference gene set captured in each new assembly (Supplementary **Table S1b** and **Fig. S1e**); and (iii) the average accuracy at the single-base level was high ($99.9996\% \pm 0.0004\%$) for the new assemblies (Supplementary **Table S1b**). In addition, the high-throughput chromosome conformation capture (Hi-C) data from four assemblies (Supplementary **Fig. S1f**) indicated the high continuity and completeness of the new assemblies. These results together suggested that the quality of all 232 new assemblies were comparable to, or better than, the levels achieved for the NIP reference genome [25], highlighting the high-quality of these new assemblies.

Next, DNA repeat sequences were annotated in the 250 high-quality genomes consisting of the 232 new assemblies and 18 existing chromosome-level assemblies downloaded from previous reports [27-31] using EDTA [32]. An average of 199.2 Mb of TEs per assembly was annotated, accounting for an average of 51.0% of the total assembly length (Supplementary **Table S1c**); overall, these values are consistent with a previous report [15]. The LTR assembly index (LAI) [33] was used to evaluate the integrity of the TE sequences in these new assemblies, and the average LAI per assembly was 21.3 ± 3.9 (Supplementary **Table S1b** and **Fig. S1g**), which

reached the “gold standard” level (LAI >20). Furthermore, comparison of the NH231 (9311) assembly sequence with bacterial artificial chromosome (BAC) clones [34] sequenced using Sanger technology showed that both sequences were of similar quality in the TE regions (Supplementary **Table S1d** and **Fig. S1h-i**).

We found that the accessions were highly diverse in terms of TE numbers. The coefficient of variation and standard deviation for the numbers of TEs per family across all 250 accessions were 0.01–0.58 and 70–3,257, respectively (**Fig. 1c**; Supplementary **Table S1c**). The variation in the total number of TEs across these 250 accessions allowed us to study the impacts of TEs on rice evolution. For example, in our previous study we found that the observed variation in genome size can be primarily explained by the number of LTRs [11]. Using our present TE datasets for *Osi*, *Osj*, and *Or* accessions, we further found that genome size variation in Asian rice was mainly due to the variation in the number of *Gypsy* elements (**Fig. 1d**; Supplementary **Fig. S1j**).

To survey the landscape of TE variation in Asian rice, we compared 250 chromosomal-level assemblies against the Nipponbare assembly using minimap2 (version 2.24-r1122) [35]. Presence/absence variants (PAVs, >50 bp) were called using custom perl scripts. When more than 90% of the sequences of a PAV are TE sequences, we consider the PAV as a TE variation (or TE-derived PAV) (Supplementary **Fig. S1k**) [36]. A pan-TE map was developed eventually (total 177,084 non-redundant TE variations, **Fig. 1b**). To quantitatively estimate the accuracy of our pan-TE map, we manually examined 200 randomly selected TE variations by visualizing the corresponding long-read mapping using the Integrative Genomics Viewer Browser. Using this method, we estimated the accuracy of our pan-TE map to be 97.5% (Supplementary **Table S1e**). We compared the TE variations identified in *Os* accessions with those previously detected using short-read sequences from 3K-RG dataset [11], and found that the average length of TE variation we identified in the 218 *Os* accessions (3,813 bp) was larger than that of in the 3K-RG dataset (346 bp) [11, 21]. Notably, almost half of TE variations were relatively long (the length of 47.3% TE variations were longer than 1 kb) (**Fig. 1e**), and 14.0% of TE variations contained at least two TE sequences. Together, these results suggested that using high-quality assemblies improved the identification of TE variations.

Inference and characterization of TE variations representing a derived state in Asian rice

In previous studies [11, 22] of population-scale TE variations, there were no rice outgroup accessions, making it hard to distinguish whether a given accession was with the ancestral state or the derived state in a given TE variation. In addition, without outgroups, it was not possible to determine whether each TE variation was the result of a TE insertion (INS) or deletion (DEL) event. Since all the three outgroups, including one *O. glaberrima* [15], one *O. barthii* [15] and one *O. glumaepatula* (<https://www.ncbi.nlm.nih.gov/nuccore/CM002522.2>), are ancestral relatives to Asian rice, and there has been no introgression between them [37, 38], we were able to use them as outgroups to represent the ancestral states of TE variations, and to infer the state of each TE variation in each accession. A TE variation, which have both derived state and ancestral state in Asian rice accessions, were defined as a derived TE variation (henceforth termed “dTE”). Derived state indicates the genotype of the TE variation in a given accession is different from outgroups, while ancestral state indicates the genotype of the TE variation in a given accession is the same as outgroups (**Fig. 1b** and **Fig. 2a**). We found 169,798 non-redundant dTEs (95.8% of non-redundant TE variations) in the Asian accessions with a total length of 647.9 Mb (**Fig. 2b**), which was about 3.5-fold higher than the total length of TE sequences in NIP (185.4 Mb). Most of the non-redundant dTE sequences were contributed by *Or* (**Fig. 2b**), possibly because of the reference bias using Nipponbare as the reference and higher genetic diversity of *Or* compared with *Osi* and *Osj*. An obviously higher number of dTEs INS events compared with DEL events was observed in the Asian rice population (**Fig. 2b**). However, except for *Or* accessions, the number/length of dTEs DELs in each accession was significant higher than that of dTEs INSs in each accession (**Fig. 2c**, $P < 0.001$). We speculate that this might be due to the unbalanced frequency between dTE INS and DEL events in Asian rice populations. 99,840 (72.9%) dTE INSs and 10,258 (31.3%) dTE DELs were present in one or only a few accessions (present frequency < 0.05) (Supplementary **Fig. S2a**). When comparing the ratio between INSs and DELs over present frequency of dTE variation, we found a decreased curve (**Fig. 2d**; Supplementary **Fig. S2a-b**), which suggests that the clearing speed to dTE INS and DEL might be different in rice evolution, with a faster clearing speed to dTE INS.

By further analyzing the dTE numbers and lengths per TE family, we found that the majority of dTEs were *Gypsy*, *Helitron*, and *Copia* elements (**Fig. 2e**), which was consistent with the high

proportion of these elements in each genome (**Fig. 1a**). The inference of dTEs can help reveal the direction and biological significance of natural and/or artificial selection acting on TEs. For example, a 977 bp dTE (*Helitron*) INS near the promoter of a transcription factor gene, *MYB61*, was reported to regulate nitrogen utilization and biomass production in rice, and the *Helitron* INS was found in *Osj* accessions, but not *Or* or *Osi* accessions or the genomes of the CC and EE outgroups in rice in previous study [38]. However, according to our high-quality Asian pan-TE map, we found the *Helitron* INS in some of the *Or* and *Osi* accessions (**Fig. 2f**), which indicated that the *Helitron* INS occurred before rice domestication. In addition, we found that the *Helitron* sequence varied among Asian accessions (Supplementary **Fig. S2c**). Notably, some *Osj* accessions had two copies of this *Helitron* INS, and the two copies were close together (~145 kb). Because the dTE contains a structurally intact *Helitron* sequence, we estimated the *Helitron* insertion time for each copy in accessions with this dTE and found that these *Helitron* sequences were relatively young overall (Supplementary **Table S1g**).

In addition, we calculated the number of dTEs nearby each gene (within ± 2 kb of a gene body) in our Asian pan-TE map and found that 31.8% of genes were conserved across all Asian accessions (no nearby dTEs were found), while almost half of genes (47.4%) contained two or more dTEs (**Fig. 2g**). Gene Ontology (GO) analyses showed that genes containing no dTEs were significantly enriched in pathways related to basic metabolism and biosynthesis of metabolites ($P < 0.05$), including cell death, photosynthesis, gene expression, amide biosynthetic process, and metabolic process (Supplementary **Fig. S2d** and **Table S1h**). In contrast, genes containing several dTEs (i.e., the number of nearby dTEs ranked in the top 5% of all genes in NIP) were significantly enriched for abiotic and biotic response genes ($P < 0.05$), such as resistance genes (*Xa1* and *Pita*) and heat shock protein genes (*OsSHSP2* and *OsHSPI*; **Fig. 2h**; Supplementary **Table S1h**), indicating that TE variations may play important roles in environmental adaptation in nature. This speculation is also supported by a significant negative correlation between latitude and the number of dTEs in each Asian rice accessions (Supplementary **Fig. S2e-f**).

Contribution of TE variations to rice domestication and differentiation

TEs can accumulate in regions of low recombination due to the action of Hill-Robertson interference [39], as we found a significant negative correlation between dTEs diversity and the

recombination rate of genome in rice (**Fig. 3a**; Supplementary **Fig. S3a**). However, we found a significantly positive correlation between *Gypsy* diversity and the recombination rate of genome in *Osi* accessions (Supplementary **Fig. S3b**), and the differences in the correlation directionality across TE types and subpopulations have been reported in other plants [9, 39], indicating the INs and DELs of TE variations are a potentially rich source of variation in the plant adaptive evolution. Numerous plant domestication and agronomic traits have been associated with particular TE variations, thus we speculated that dTEs may play an important role in evolutionary processes, such as domestication and differentiation in rice. To explore the impacts of TE variations on the domestication and differentiation of Asian rice, we investigated TE variations in 247 diverse Asian rice accessions. A phylogenetic tree constructed using our dTEs dataset for the Asian rice accessions clearly divided the accessions into the groups of *Osi*, *Osj*, *Aus*, and *Or* (Supplementary **Fig. S3c**), consistent with the SNP-based rice phylogeny (**Fig. 1a**). Meanwhile, the results of admixture (Supplementary **Fig. S3d**) and PCA (Supplementary **Fig. S3e-f**) using the dTEs dataset also revealed that TE variations were tightly associated with Asian rice domestication and differentiation.

Next, we calculated the average number of dTEs in the selective windows (i.e., ranked in the top 5% of 100kb F_{ST} windows for SNPs) between subpopulations. dTEs were significantly enriched in selective windows than expected at random in comparisons between *Or* and *Osi*, *Or* and *Osj*, and *Osi* and *Osj*, even under the conditions that we controlled the sample size of each subpopulation to the same level (**Fig. 3b**, $P < 2.2e-16$; Supplementary **Fig. S3g**), indicating that dTEs probably participated in rice domestication (from *Or* to *Osi* and *Or* to *Osj*) and differentiation (between *Osi* and *Osj*). Thus, to determine whether the observed differential distribution of dTEs genotypes among subpopulations was due to selection, we estimated the divergence of each dTE across the genome using F_{ST} analysis, and 2,141 and 2,043 of dTEs showing signatures of selection (henceforth termed “selected dTEs”; i.e., locus showing signatures of selection, i.e., F_{ST} outlier between *Or* and *Osi* [outlier = 0.85], *Or* and *Osj* [outlier = 0.84]) were detected in the *Or-Osi* and *Or-Osj* comparisons, respectively (**Fig. 3c**). In total, 3,935 selected dTEs possibly involved in the domestication processes were ultimately identified (**Fig. 3c**). Using the same F_{ST} analysis approach, we also identified 2,108 selected dTEs possibly involved in differentiation between *Osi* and *Osj* (outlier F_{ST} value = 0.94) (**Fig. 3c**). Almost all LTR types and

two DNA types, MITE and *Helitron*, were significantly enriched in rice domestication and differentiation processes ($P < 0.05$, **Fig. 3d**), while SINE and LINE were significantly enriched in the domestication processes from *Or* to *Osi* ($P < 0.05$, **Fig. 3d**).

We analyzed the genes overlapping with selected dTEs and identified 3,860 potential candidate genes (henceforth termed “dTE-genes”), including 2,992 dTE-genes with evidence of selection between *Or* and *Os* (1,634 dTE-genes between *Or* and *Osi*, and 1,580 dTE-genes between *Or* and *Osj*) and 1,750 dTE-genes with evidence of selection between *Osi* and *Osj*. Of these, we showed the selection and expression differences for 169 genes with known functions (**Fig. 3e**; Supplementary **Fig. S3h**). We found selected dTEs nearby previously identified putative domestication genes, such as *An-2* [40]. In addition, we found a selected dTE (345 bp) INS in the intron of a known gene, *OsAKR2*, between *Or* and *Osj*, which was located in a region in strong linkage disequilibrium (LD) with the known domestication gene *qSh1* (Supplementary **Fig. S3i**), we can infer this dTE INS may be also strongly selected in the domestication process due to strong linkage disequilibrium with the causal SNP variant in *qSh1*. We also found a 112 bp selected dTE (LINE) DEL in the 3'UTR of the drought tolerant gene *OsCML4* in upland *Osj* accessions, and the expression level of *OsCML4* in the upland *Osj* accessions with the dTE DEL was significantly higher than that of in the *Or* accessions without this dTE DEL (Supplementary **Fig. S3j**). Given that increased *OsCML4* expression likely causes the upland *Osj* accessions to have stronger drought resistance than the other *Os* accessions.

We found that two dTE-genes that might be involved in environmental adaptation, promoting the formation of diverse crops during the differentiation of *Osi* and *Osj*. For example, a selected dTE occurred between *Osi* and *Osj* subpopulations ($F_{ST} = 0.94$, rank: 4.42%). This 157 bp sequence with a *Helitron* was located downstream of a known heat stress gene, *OsClpB-m* [41]. It was found in the outgroups (ancestral state), some of the *Or* accessions (18 out of 27; the genotype of one accession was uncertain), and most of the *Osi* accessions (120 out of 146; the genotypes of 21 accessions were uncertain), but absent in most of the *Osj* accessions (derived state; 54 out of 61; the genotypes of remaining 7 accessions were uncertain) (Supplementary **Fig. S3k**). The *Osj* accessions with the dTE DEL event had significantly lower expression levels of *OsClpB-m* [41] (Supplementary **Fig. S3k**, $P < 0.01$) and heat tolerance (Supplementary **Fig. S3k**, $P < 0.001$) than the *Osi* accessions without the dTE DEL event.

Another dTE-gene example was the rice cold-induced *LIP19* [42] gene, which encodes a 148-amino-acid basic region/leucine zipper protein and is highly expressed in the mesophyll cells surrounding vascular bundles of rice plants. We detected a 150 bp selected dTE (*Tourist* MITE, $F_{ST} = 0.95$, rank: 3.71%) inserted in the promoter of *LIP19* (**Fig. 3f**). The 150 bp region was present in almost all *Osj* accessions (60 out of 61) but few *Osi* accessions (4 out of 146, **Fig. 3f**). The expression level of *LIP19* in the *Osj* accessions with the dTE INS was shown to be significantly higher than those *Osi* accessions without the dTE INS (**Fig. 3g**, $P < 0.001$). It is therefore possible that this dTE INS have contributed to cold tolerance in the *Osj* accessions rather than those *Osi* accessions without the dTE INS (**Fig. 3h-i**, $P < 0.001$). *Osj* accessions are distributed in the regions with a higher latitude and colder temperature, and the dTE state (including derived state and ancestral state) of the temperature adaptation genes also distributed according to the latitude (**Fig. 3j**; Supplementary **Fig. S31**), which may be affected by climate adaptability selection due to temperature. The phylogenetic tree of the dTE and *LIP19* sequences from the 247 Asian rice accessions and 3 outgroups showed that the outgroups were without the 150 bp dTE region. The *Osj* accessions and *Osi* accessions were grouped with different *Or* accessions, which was consistent with the distribution of the dTE INS event. The dTE in the *Osj* subpopulation was derived from *Or* accessions (**Fig. 3k**). The fact that the distribution of this dTE INS event was nearly fixed in the *Osj* subpopulation (60 out of 61) and nearly totally absent in the *Osi* subpopulation (4 out of 146), demonstrated that the dTE was selected during the domestication of *Osj* and *Osi*, potentially because of its effect on cold response through the modulating expression of the *LIP19* gene. These results suggested that dTEs was probably contribute to climate adaptability by altering gene expression in rice.

TE variations associated with gene expression

In many organisms, including plants and animals, TEs have been shown to be associated with differences in expression level [43]. To assess more systematically whether TEs are a potentially important source of gene expression variation [5, 12, 22], we first measured the LD of dTE with nearby SNPs and insertion-deletion polymorphisms (InDels). We observed that a large fraction of dTEs are in strong LD ($R^2 > 0.7$) with adjacent (50 kb on either side) SNPs and InDels (43.4% and 47.6%, respectively, **Fig. 4a**), and a small proportion (7.5% and 11.3%, respectively) are in

complete LD ($R^2 = 1$). However, nearly one-fourth of dTEs are in weak LD (25.2% and 30.3%, respectively, $R^2 < 0.1$), indicating that TE variations are not perfectly tagged by SNPs and InDel markers. We next calculated the number of dTEs in terms of relative position to genes and found that 79.5% of dTEs overlapping with genic regions (within ± 2 kb of a gene body) (**Fig. 4b**). Compared with the distribution of TEs in the NIP genome [25], the dTEs from the Asian pan-TE map were significantly enriched in transcriptional regulatory regions ($P < 0.05$) and depleted in coding sequences (CDSs, $P < 0.001$) and intronic regions ($P < 0.001$, **Fig. 4b**). All TE families showed a pattern similar to that of the whole dTE dataset (**Fig. 4b**), indicating that TEs may play a role in regulating gene expression.

To estimate the relative contribution of TE variations and SNPs to gene expression changes, we associated dTEs and SNPs with gene expression levels among *Os* accessions, respectively and identified expression quantitative trait loci (eQTLs). Using leaf transcriptome data from our previous study [15], we identified 3,868 genes whose expression was significantly associated with dTE (referred to as eGenes, **Fig. 4c**). Of these eGenes, 2,622 were also identified in the SNP-eQTL analysis. For example, a dTE (DTM) INS present in the intron of a shared gene *OsGluRS* [44] was found in 60 out of 61 *Osj* accessions, while the leading SNP variant with *OsGluRS* [44] gene expression variations was ~ 4.8 kb away from the gene (Supplementary **Fig. S4a**), indicating that the dTE is potentially a crucial variation altering *OsGluRS* [44] expression. Knockdown lines of *OsGluRS* [44] in *Osj* accessions resulted in an abnormal phenotype characterized by yellow-green leaves when plants were grown at 26°C or higher (nonpermissive temperature) [44]. We also found similar results at the population level; accessions with the dTE INS (the derived state) showed significantly lower expression and lower heat-resistance than those without the dTE INS (the ancestral state, Supplementary **Fig. S4a**). Another example is a known positive regulator gene of bacterial blight resistance, *OsRPIL1* [45]. A 1.0 kb dTE (*Helitron*) INS in the gene promoter was identified in most *Osj* accessions (57 out of 61) and a few *Osi* accessions (13 out of 146) (Supplementary **Fig. S4b**). The expression of *OsRPIL1* was previously shown to ensure appropriate bacterial blight resistance in rice [45]. We found that the accessions with the dTE INS had significantly higher expression levels and possibly associated with lower susceptibility to bacterial blight than the accessions without the dTE INS (Supplementary **Fig. S4b**), indicating that this dTE may play a role in increasing bacterial blight resistance. For the 2,622 shared eGenes

between TE-eQTL and SNP-eQTL analysis, we compared the expression variance explained by the most significantly associated dTEs and SNPs. For ~48.9% of these genes, the leading associated dTEs could explain more expression variance than the leading associated SNPs.

Importantly, the expression levels of 1,246 eGenes were only associated with dTEs and not associated with SNP data (**Fig. 4c**), demonstrating the importance of considering dTEs in gene expression. For example, the expression of a rice NADPH oxidase gene, *OsRbohB* [46], was significantly associated with a 328 bp dTE (DNA TE PILE) INS event in its promoter region (**Fig. 4d**). We did not find any significant association of *OsRbohB* expression with SNPs. The dTE INS was associated with decreased expression of *OsRbohB* in the accessions with the dTE INS (**Fig. 4e**), and the accessions with the dTE INS had reduced thousand grain weight compared with those without the dTE INS (**Fig. 4f**) in both the *Osi* and *Osj* accessions (Supplementary **Fig. S4c**). It seems plausible that this dTE INS may be contributed to thousand grain weight, consistent with the phenotype of the *OsRbohB* mutant showing lower thousand grain weight in previous study [46]. Compared to the one of accessions (such as NH016) without the dTE INS, the one (such as NH243) of accessions with the dTE INS in a *OsRbohB* promoter fragment significantly reduced its relative expression activity (**Fig. 4g**). These results validated the incomplete LD between dTEs and small variants (SNPs and InDels) and highlighted the potential of dTEs to explain additional variation in gene expression and agronomic traits.

TE variations associated with agronomic traits

To demonstrate how the pan-TE map can be used to facilitate the identification of agronomic traits associated with TE variations, we performed a genome-wide association study (GWAS) of agronomic traits in *Oryza* accessions. A *GW5* peak was previously detected in a GWAS on grain width using SNPs or TE variations genotyped using short-read data [47]. We also detected the same signal for grain width. In addition, we identified a new locus on chromosome 7 harboring a 131 bp dTE (*Stowaway* MITE) INS that was significantly associated with grain width (Supplementary **Fig. S5a**). The locus was not identified using SNPs or the previous TE variations based on short-reads data. Rice accessions with the dTE INS (derived state) was shown to be significantly lower grain width than those without the dTE INS (ancestral state) ($P < 0.001$, Supplementary **Fig. S5b**). Within 5 kb region of the TE peak, there were two candidate genes. One

of them was a gene, *LOC_Os07g41000* [48], previously reported to regulate grain width (Supplementary **Fig. S5b-c**), indicating that the dTE might control grain width by influencing *LOC_Os07g41000* [48].

Using phenotypes related to heat tolerance to conduct TE-GWAS, we identified two dTEs peak loci strongly associated with heat stress in *Osi* accessions; these loci were not identified using SNP dataset of *Osi* accessions (Supplementary **Fig. S5d-g**). From the candidate genes near each dTEs peak (5kb on either side), we found a 263 bp dTE (*Gypsy*) DEL, which was associated with the level of heat resistance, was located near an OsFBX121 F-box domain gene, *LOC_Os04g11890* (Supplementary **Fig. S5d-e**, $P < 0.001$). F-box proteins play important roles in responses to environmental stimuli [49], indicating that the dTE might affect heat resistance by affecting *LOC_Os04g11890*. Likewise, another TE locus associated with leaf survival rate under heat stress was located near a calcium-binding protein gene, *LOC_Os03g07470* (rice annexin *OsANN1*) [50], which could potentially confer enhances heat stress tolerance by modulating the production of H_2O_2 (Supplementary **Fig. S5f-g**, $P < 0.01$). Together, these results indicated that the two dTEs are associated with heat stress.

We also found that a dTE peak harboring a 1.0 kb dTE (*Gypsy*) DEL on chromosome 6 was strongly associated with seed setting rate under cold stress, and this locus was not identified based on SNP data (**Fig. 4h**; Supplementary **Fig. S5h**). This dTE DEL was located 1.6 kb upstream of the bolA family gene *LOC_Os06g28970* [51], which has been linked to stress response pathways. The absence or overexpression of bolA proteins could modulate the response to a variety of environmental challenges [51]. A follow-up analysis showed that the seed setting rate in accessions with the dTE DEL was significantly higher than that in accessions without the dTE DEL (**Fig. 4i**, $P < 0.01$). The expression level of *LOC_Os06g28970* in accessions with the dTE DEL was significantly higher than that in accessions without the dTE DEL (**Fig. 4j**, $P < 0.05$). To identify the high-confidence variants predicted to be associated with seed setting rate under cold stress, we performed Sum of Single Effects (SuSiE) [52] analysis to fine-map the dominance loci ($PIP = 1$) using dTE and SNP variants located near (100 kb on either side) the peak. Then combining these fine-mapping results with the significant TE-GWAS and SNP-GWAS peaks, we confirmed that the 1.0kb dTE DEL associated with *LOC_Os06g28970* [51] was most likely to be the causal variant affecting seed setting rate under cold stress. These results allowed us to infer

that dTE might make widespread contributions to rice complex traits.

DISCUSSION

Rice is a major food crop for billions of people globally [17], and obtaining a deeper understanding of rice phenotypic variation and crop domestication can significantly benefit agriculture, world food security, and the biological and genomic research communities. In the present study, we took advantage of 15 existing and 232 new high-quality chromosomal-level assemblies of diverse accessions to generate an Asian rice pan-TE map. Subsequently, using three non-Asian rice accessions as outgroups, we used the pan-TE map to infer TE variations representing the derived state, and analyzed their characteristics in Asian rice accessions. Compared to SNPs, TE variations often explain more phenotypic variance and can be used for genomic prediction [53]. The reason for this could be that, as compared with SNPs, TE variations are more frequently the causal mutation. For example, by performing dTE-eQTL analysis, we identified 1,246 genes whose expression was significantly associated with dTEs but not identified based on SNP data. Additionally, using dTE data to conduct GWAS of agronomic traits in rice, we successfully uncovered multiple novel associations that could not be detected using SNP data (Fig. 4h).

Given that knowing the ancestral or derived state of each TE can support inferences about the evolutionary relationships between rice accessions and the genetic basis of phenotypic diversity in the rice populations, these rich dTE datasets represent important resources for investigations of rice evolution and domestication history in functional genomics studies. Using the pan-TE map, we found that the number of dTE INS events was greater than dTE DEL events. TE variation frequency is commonly used to mirror TE age, because low frequency TE variations are much younger than those in the outgroup [54]. Consistent with this assumption, the vast majority of low frequency dTE INSs were younger and more quickly removed than dTE DELs in nature, implying that these sequences might be deleterious and selected during in artificial domestication or natural selection [55].

The construction of a species' pan-genome enables us to resolve hidden genomic complexity involving TE variations, and re-evaluate their role in genome evolution and generation of phenotypic variation. Previous studies utilized Asian cultivated rice, such 208 *Os* accessions [22]

and 3010 *Os* accessions[11] and 176 *Os* accessions [56], based on the short-read data to identify TE and develop TE dataset, without *Or* accessions, while our pan-TE variation map included 218 *Os* and 29 *Or* accessions. We can analyze the changes in TE variation during rice domestication processes, and found 3,935 selected dTEs seem likely to be involved in the domestication process from *Or* to *Osi* or *Osj*. In addition, our use of high-quality chromosome-level genome assemblies enabled us to identify complex TE variation. We found dTEs that contain two or more TEs in our pan-TE map, such as a dTE consisting of one *Gypsy*, one *Helitron*, and one *Copia* element, which are also likely to contribute to phenotypic diversity. Using the high-quality TE dataset, we could identify TEs contributions to environmental adaptation and domestication, and some rare TE mutations may have greater potential for gene cloning and molecular breeding [57].

In addition, our reference-quality genomes will be helpful to the very large number of research groups doing various basic biological studies [58], such as those developing genome editing technologies and promoting successful mapping of causal mutations positioned within highly complex and/or duplicated regions. The Asian pan-TE map also offers a platform that enables researchers to obtain substantially more information than using a few accessions.

MATERIALS AND METHODS

Detailed materials and methods are available in the supplementary information.

DATA AVAILABILITY

All new assemblies of 232 accessions in the present study have been deposited at the Genome Warehouse (<https://bigd.big.ac.cn/gwh/>) under PRJCA018476. The TE annotation of all assemblies have been deposited at the Figshare (https://figshare.com/articles/dataset/Annotations_of_250_high_quality_rice_genomes/23803956).

ACKNOWLEDGEMENTS

We are very grateful to the High-performance computing centers of Agricultural Genomics Institute at Shenzhen, and computing centers of China National Rice Research Institute for their technical support.

FUNDING

This research was supported by the National Natural Science Foundation of China (32188102, 32101718), Guangdong Basic and Applied Basic Research Foundation (2023B1515020053), Innovation Program of Chinese Academy of Agricultural Sciences, and the Youth innovation of Chinese Academy of Agricultural Sciences (Y2023QC36).

AUTHOR CONTRIBUTIONS

L.S., Q.Q., and Y.Zhou conceived the idea and designed the project; X.Li., X.D., H.H., and Y.Lv collected and downloaded the samples and investigated sequencing data; X.Li participated in the execution of sequence assembly, quality assessment and TE annotation; X.D. accomplished the construction and analysis of pan-TE; H.H. fulfilled the inferring of derived TE; Y.Lv completed the analysis of the validation of TE; L.Y. performed the SNP and GWAS analysis; W.C.H was responsible for FST analysis; C.L. and B.Z were responsible for TE families; H.Wei and Q.Z. did the molecular biology experiments; X.W. and Y.Leng were responsible for Hi-C data analysis; X.Yu. and Q.Yuan implemented the construction of TE phylogenetic tree; H.Z and X.P.L performed eQTL analysis; B.T.Z and D.Chen completed the analysis of TE haplotype; X.Yu. and X.L.Cao accomplished the construction of SNP phylogenetic tree; C.S. and T.W. were responsible for estimation of recombination rate; W.Chen and M.G. implemented the calculate of TE insertion time; H.Q. and Y.Cui drew haplotype diagram; Q.X and Z.Z. did the enrichment and Fisher's exact test across different family; L.S., Q.Q. and Y.Zhou coordinated all co-authors and revised the manuscript.

Competing interest. The authors declare that they no conflict of interest.

REFERENCES

1. Feschotte C, Jiang N, Wessler SR. Plant transposable elements: where genetics meets genomics. *Nat Rev Genet.* 2002; **3**(5): 329-341. doi: 10.1038/nrg793
2. Mc CB. Chromosome organization and genic expression. *Cold Spring Harb Symp Quant Biol.* 1951; **16**: 13-47. doi: 10.1101/sqb.1951.016.01.004
3. Wicker T, Sabot F, Hua-Van A *et al.* A unified classification system for eukaryotic transposable elements. *Nat Rev Genet.* 2007; **8**(12): 973-982. doi: 10.1038/nrg2165
4. Chénais B. Transposable Elements and Human Diseases: Mechanisms and Implication in the Response to Environmental Pollutants. *Int J Mol Sci.* 2022; **23**(5). doi:

10.3390/ijms23052551

5. Cai X, Lin R, Liang J *et al.* Transposable element insertion: a hidden major source of domesticated phenotypic variation in *Brassica rapa*. *Plant Biotechnol J.* 2022; **20**(7): 1298-1310. doi: 10.1111/pbi.13807
6. Niu C, Jiang L, Cao F *et al.* Methylation of a MITE insertion in the *MdRFNR1-1* promoter is positively associated with its allelic expression in apple in response to drought stress. *Plant Cell.* 2022; **34**(10): 3983-4006. doi: 10.1093/plcell/koac220
7. Domínguez M, Dugas E, Benchouaia M *et al.* The impact of transposable elements on tomato diversity. *Nat Commun.* 2020; **11**(1): 4058. doi: 10.1038/s41467-020-17874-2
8. Zhang L, Hu J, Han X *et al.* A high-quality apple genome assembly reveals the association of a retrotransposon and red fruit colour. *Nat Commun.* 2019; **10**(1): 1494. doi: 10.1038/s41467-019-09518-x
9. Lisch D. How important are transposons for plant evolution? *Nat Rev Genet.* 2013; **14**(1): 49-61. doi: 10.1038/nrg3374
10. Studer A, Zhao Q, Ross-Ibarra J *et al.* Identification of a functional transposon insertion in the maize domestication gene *tb1*. *Nat Genet.* 2011; **43**(11): 1160-1163. doi: 10.1038/ng.942
11. Carpentier MC, Manfroi E, Wei FJ *et al.* Retrotranspositional landscape of Asian rice revealed by 3000 genomes. *Nat Commun.* 2019; **10**(1): 24. doi: 10.1038/s41467-018-07974-5
12. Wei KH, Mai D, Chatla K *et al.* Dynamics and Impacts of Transposable Element Proliferation in the *Drosophila nasuta* Species Group Radiation. *Mol Biol Evol.* 2022; **39**(5). doi: 10.1093/molbev/msac080
13. Wing RA, Purugganan MD, Zhang Q. The rice genome revolution: from an ancient grain to Green Super Rice. *Nat Rev Genet.* 2018; **19**(8): 505-517. doi: 10.1038/s41576-018-0024-z
14. Qin P, Lu H, Du H *et al.* Pan-genome analysis of 33 genetically diverse rice accessions reveals hidden genomic variations. *Cell.* 2021; **184**(13): 3542-3558 e3516. doi: 10.1016/j.cell.2021.04.046
15. Shang L, Li X, He H *et al.* A super pan-genomic landscape of rice. *Cell Res.* 2022; **32**(10): 878-896. doi: 10.1038/s41422-022-00685-z
16. Tao Y, Zhao X, Mace E *et al.* Exploring and Exploiting Pan-genomics for Crop Improvement. *Mol Plant.* 2019; **12**(2): 156-169. doi: 10.1016/j.molp.2018.12.016
17. Wang W, Mauleon R, Hu Z *et al.* Genomic variation in 3,010 diverse accessions of Asian cultivated rice. *Nature.* 2018; **557**(7703): 43-49. doi: 10.1038/s41586-018-0063-9
18. Gao Y, Xu Z, Zhang L *et al.* *MYB61* is regulated by *GRF4* and promotes nitrogen utilization and biomass production in rice. *Nat Commun.* 2020; **11**(1): 5219. doi: 10.1038/s41467-020-19019-x
19. Wu B, Meng J, Liu H *et al.* Suppressing a phosphohydrolase of cytokinin nucleotide enhances grain yield in rice. *Nat Genet.* 2023. doi: 10.1038/s41588-023-01454-3
20. Mao D, Tao S, Li X *et al.* The Harbinger transposon-derived gene PANDA epigenetically

- coordinates panicle number and grain size in rice. *Plant Biotechnol J*. 2022; **20**(6): 1154-1166. doi: 10.1111/pbi.13799
21. Fuentes RR, Chebotarov D, Duitama J *et al*. Structural variants in 3000 rice genomes. *Genome Res*. 2019; **29**(5): 870-880. doi: 10.1101/gr.241240.118
 22. Castanera R, Morales-Díaz N, Gupta S *et al*. Transposons are important contributors to gene expression variability under selection in rice populations. *Elife*. 2023; **12**. doi: 10.7554/eLife.86324
 23. Hu, J. *et al*. An efficient error correction and accurate assembly tool for noisy long reads. *bioRxiv* 2023.03.09.531669 (2023) doi:10.1101/2023.03.09.531669.
 24. Hu J, Fan J, Sun Z *et al*. NextPolish: a fast and efficient genome polishing tool for long-read assembly. *Bioinformatics*. 2020; **36**(7): 2253-2255. doi: 10.1093/bioinformatics/btz891
 25. Kawahara Y, de la Bastide M, Hamilton JP *et al*. Improvement of the *Oryza sativa* Nipponbare reference genome using next generation sequence and optical map data. *Rice (N Y)*. 2013; **6**(1): 4. doi: 10.1186/1939-8433-6-4
 26. Simão FA, Waterhouse RM, Ioannidis P *et al*. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*. 2015; **31**(19): 3210-3212. doi: 10.1093/bioinformatics/btv351
 27. Zhou Y, Chebotarov D, Kudrna D *et al*. A platinum standard pan-genome resource that represents the population structure of Asian rice. *Sci Data*. 2020; **7**(1): 113. doi: 10.1038/s41597-020-0438-2
 28. Song JM, Xie WZ, Wang S *et al*. Two gap-free reference genomes and a global view of the centromere architecture in rice. *Mol Plant*. 2021; **14**(10): 1757-1767. doi: 10.1016/j.molp.2021.06.018
 29. Dong PF, Tu XY, Chu PY *et al*. 3D Chromatin Architecture of Large Plant Genomes Determined by Local A/B Compartments. *Mol Plant*. 2017; **10**(12): 1497-1509. doi: 10.1016/j.molp.2017.11.005
 30. Xie X, Du H, Tang H *et al*. A chromosome-level genome assembly of the wild rice *Oryza rufipogon* facilitates tracing the origins of Asian cultivated rice. *Sci China Life Sci*. 2021; **64**(2): 282-293. doi: 10.1007/s11427-020-1738-x
 31. Zhang H, Wang Y, Deng C *et al*. High-quality genome assembly of Huazhan and Tianfeng, the parents of an elite rice hybrid Tian-you-hua-zhan. *Sci China Life Sci*. 2022; **65**(2): 398-411. doi: 10.1007/s11427-020-1940-9
 32. Ou S, Su W, Liao Y *et al*. Benchmarking transposable element annotation methods for creation of a streamlined, comprehensive pipeline. *Genome Biol*. 2019; **20**(1): 275. doi: 10.1186/s13059-019-1905-y
 33. Ou S, Chen J, Jiang N. Assessing genome assembly quality using the LTR Assembly Index (LAI). *Nucleic Acids Res*. 2018; **46**(21): e126. doi: 10.1093/nar/gky730
 34. Pan Y, Deng Y, Lin H *et al*. Comparative BAC-based physical mapping of *Oryza sativa* ssp.

- indica var. 93-11 and evaluation of the two rice reference sequence assemblies. *Plant J.* 2014; **77**(5): 795-805. doi: 10.1111/tpj.12412
35. Li H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics.* 2018; **34**(18): 3094-3100. doi: 10.1093/bioinformatics/bty191
36. Chen J, Liu Y, Liu M *et al.* Pangenome analysis reveals genomic variations associated with domestication traits in broomcorn millet. *Nat Genet.* 2023; **55**(12): 2243-2254. doi: 10.1038/s41588-023-01571-z
37. Choi JY, Platts AE, Fuller DQ *et al.* The Rice Paradox: Multiple Origins but Single Domestication in Asian Rice. *Mol Biol Evol.* 2017; **34**(4): 969-979. doi: 10.1093/molbev/msx049
38. Stein JC, Yu Y, Copetti D *et al.* Genomes of 13 domesticated and wild rice relatives highlight genetic conservation, turnover and innovation across the genus *Oryza*. *Nat Genet.* 2018; **50**(2): 285-296. doi: 10.1038/s41588-018-0040-0
39. Dolgin ES, Charlesworth B. The effects of recombination rate on the distribution and abundance of transposable elements. *Genetics.* 2008; **178**(4): 2169-2177. doi: 10.1534/genetics.107.082743
40. Gu B, Zhou T, Luo J *et al.* An-2 Encodes a Cytokinin Synthesis Enzyme that Regulates Awn Length and Grain Production in Rice. *Mol Plant.* 2015; **8**(11): 1635-1650. doi: 10.1016/j.molp.2015.08.001
41. Singh A, Singh U, Mittal D *et al.* Genome-wide analysis of rice ClpB/HSP100, ClpC and ClpD genes. *BMC Genomics.* 2010; **11**: 95. doi: 10.1186/1471-2164-11-95
42. Shimizu H, Sato K, Berberich T *et al.* LIP19, a Basic Region Leucine Zipper Protein, is a Fos-like Molecular Switch in the Cold Signaling of Rice Plants. *Plant and Cell Physiology.* 2005; **46**(10): 1623-1634. doi: 10.1093/pcp/pci178
43. Cridland JM, Thornton KR, Long AD. Gene expression variation in *Drosophila melanogaster* due to rare transposable element insertion alleles of large effect. *Genetics.* 2015; **199**(1): 85-93. doi: 10.1534/genetics.114.170837
44. Liu W, Fu Y, Hu G *et al.* Identification and fine mapping of a thermo-sensitive chlorophyll deficient mutant in rice (*Oryza sativa* L.). *Planta.* 2007; **226**(3): 785-795. doi: 10.1007/s00425-007-0525-z
45. Wang X, Chen J, Yang Y *et al.* Characterization of a Novel NBS-LRR Gene Involved in Bacterial Blight Resistance in Rice. *Plant Mol Biol Rep.* 2013; **31**(3): 649-656. doi: 10.1007/s11105-012-0537-0
46. Shi Y, Chang YL, Wu HT *et al.* OsRbohB-mediated ROS production plays a crucial role in drought stress tolerance of rice. *Plant Cell Rep.* 2020; **39**(12): 1767-1784. doi: 10.1007/s00299-020-02603-2
47. Akakpo R, Carpentier MC, Ie Hsing Y *et al.* The impact of transposable elements on the structure, evolution and function of the rice genome. *New Phytol.* 2020; **226**(1): 44-49. doi:

10.1111/nph.16356

48. Misra G, Badoni S, Anacleto R *et al.* Whole genome sequencing-based association study to unravel genetic architecture of cooked grain width and length traits in rice. *Scientific Reports*. 2017; **7**(1): 12478. doi: 10.1038/s41598-017-12778-6
49. Cao Y, Yang Y, Zhang H *et al.* Overexpression of a rice defense-related F-box protein gene *OsDRF1* in tobacco improves disease resistance through potentiation of defense gene expression. *Physiol Plant*. 2008; **134**(3): 440-452. doi: 10.1111/j.1399-3054.2008.01149.x
50. Qiao B, Zhang Q, Liu D *et al.* A calcium-binding protein, rice annexin *OsANN1*, enhances heat stress tolerance by modulating the production of H₂O₂. *J Exp Bot*. 2015; **66**(19): 5853-5866. doi: 10.1093/jxb/erv294
51. Fleurie A, Zoued A, Alvarez L *et al.* A *Vibrio cholerae* BolA-Like Protein Is Required for Proper Cell Shape and Cell Envelope Integrity. *mBio*. 2019; **10**(4). doi: 10.1128/mBio.00790-19
52. Wang G, Sarkar A, Carbonetto P *et al.* A simple new approach to variable selection in regression, with application to genetic fine mapping. *J R Stat Soc Series B Stat Methodol*. 2020; **82**(5): 1273-1300. doi: 10.1111/rssb.12388
53. Vourlaki IT, Castanera R, Ramos-Onsins SE *et al.* Transposable element polymorphisms improve prediction of complex agronomic traits in rice. *Theor Appl Genet*. 2022; **135**(9): 3211-3222. doi: 10.1007/s00122-022-04180-2
54. Juan J, Yong-Chao X, Zhi-Qin Z *et al.* Forces driving transposable element load variation during *Arabidopsis* range expansion. *bioRxiv*. 2022: 2022.2012.2028.522087. doi: 10.1101/2022.12.28.522087
55. Bergman CM, Quesneville H, Anxolabéhère D *et al.* Recurrent insertion and duplication generate networks of transposable element sequences in the *Drosophila melanogaster* genome. *Genome Biol*. 2006; **7**(11): R112. doi: 10.1186/gb-2006-7-11-r112
56. Yan H, Haak DC, Li S *et al.* Exploring transposable element-based markers to identify allelic variations underlying agronomic traits in rice. *Plant Commun*. 2022; **3**(3): 100270. doi: 10.1016/j.xplc.2021.100270
57. Wang T, He W, Li X, *et al.* A rice variation map derived from 10,548 rice accessions reveals the importance of rare variants. *Nucleic Acids Res*. 2023; **51**(20):10924-10933. doi:10.1093/nar/gkad840
58. Shang L, He W, Wang T, *et al.* A complete assembly of the rice Nipponbare reference genome. *Mol Plant*. 2023 ;**16** (8):1232-1236. doi:10.1016/j.molp.2023.08.003




Fig. 1 Construction of a pan-TE map and evaluation of TE variations in Asian accessions using chromosome-level genomes.

a Landscape of genome size and TE content across different subpopulations. Phylogeny of 250 accessions were based on whole-genome SNPs (top); accessions in different subpopulations are indicated by different colors. *Osi*, *Aus*, *Osj*, *Or*, and outgroup respectively refer to *O. sativa indica*, *O. sativa aus*, *O. sativa japonica*, *O. rufipogon*, and three non-Asian accessions (one *O. glaberrima*, one *O. barthii*, and one *O. glumaepatula*). Length of genome and TE content (Mb) in each genome were indicated (bottom). The length of *Gypsy*, *Copia*, DTC, DTA, DTT, DTM, DTH, *Helitron*, LINE, and SINE elements in each genome are indicated. **b** Overview of the pipeline for pan-TE map construction. Firstly, 232 high-quality chromosomal-level assemblies were de novo assembled by integrating public long-read and short-read data. After combining the de novo assemblies with 18 existing assemblies, the TE sequences were annotated and a

TE library was generated. To construct a pan-TE map, the TE variations were identified by combining the results of genome alignment, the TE library, and long read data. Subsequently three non-Asian accessions were used as outgroups (henceforth termed “outgroup”) to infer whether a given TE variation was in derived state or ancestral state in each accession. A TE variation, which have both derived state and ancestral state in Asian rice accessions, was defined as derived TE variation (henceforth termed “dTE”). Ancestral state indicates that the genotype of the locus in a given accession (0/0) is the same as that of the outgroup (0/0); derived state indicates that the genotype of the locus in a given accession (1/1 or 0/1) is different from that of the outgroup (0/0), including homozygous (1/1) and heterozygous (0/1) genotype. Finally, a dTE genotype dataset in matrix format is generated for use in downstream analysis, including domestication, gene expression, and GWAS. **c** The copy number variation for different TE families in 250 natural accessions. X axis represents the copy number variation for each TE family across accessions, evaluated as coefficient of variation (CV); y axis represents the average number of TEs in each family; z axis represents the differences in total TE number for each family among accessions in total TE number, evaluated as standard deviation (SD). **d** Pearson correlation coefficients for comparisons between total length of *Gypsy* elements and genome size across different subpopulations. Colored dots and lines indicate data from each subpopulation. **e** Length distributions of TE variations in the non-redundant TE dataset for Asian accessions.




Fig. 2 Inference and characterization of TE variations representing a derived state across Asian accessions.

a The landscape of dTEs across the Asian accessions. Three non-Asian accessions were used as outgroups (henceforth termed “outgroup”) to infer whether a given TE variation was in derived state or ancestral state in each accession. The TE variations, which have both derived state and ancestral state in Asian rice accessions, were defined as derived TE variations (henceforth termed “dTEs”). Ancestral state (green) indicates that the genotype of a TE variation in an accession is consistent with that of the outgroup; derived state indicates that the genotype of a TE variation, homozygous (pink) or heterozygous (purple), in an accession is different from that of the outgroup. Gray boxes

indicate that the genotype is uncertain (NA). *Osi*, *Aus*, *Osj*, *Or*, and outgroup respectively refer to *O. sativa indica*, *O. sativa aus*, *O. sativa japonica*, *O. rufipogon*, and three non-Asian accessions (*O. glaberrima*, *O. barthii*, and *O. glumaepatula*). **b** Total length and number of non-redundant dTEs and sequences detected across different subpopulations. DEL and INS refer to deletion and insertion events identified by comparing the Asian accessions to the outgroups, respectively. “All” refers to all the Asian rice accessions in the present study. **c** Average length and number of dTEs for each accession across different subpopulations. *** $P < 0.001$, Significance was determined using the Student's *t*-test. **d** The count ratio of INS events to DEL events for different frequencies of dTEs in Asian rice accessions. **e** Total length and number of dTEs for TE families in Asian rice accessions. **f** A *Helitron* in the *MYB61* promoter region was found in all *Osj* accessions in a previous study, but was undetectable in *Or* and *Osi* accessions and the rice outgroup genomes CC and EE. However, the *Helitron* inserted in some *Or* and *Osi* accessions in our high-quality Asian pan-TE map. **g** Distribution of the number of dTEs near each gene (within ± 2 kb of a gene body). **h** Gene ontology (GO) analyses of the genes containing several dTEs (“the genes” were ranked by the number of dTEs overlapping with their genic region, and the top 5% of ranked genes were used in the GO analyses). BP, CC, and MF respectively refer to biological process, cellular component, and molecular function.




Fig. 3 Contribution of dTEs to rice domestication and differentiation.

a Pearson correlation coefficient between dTE average pairwise diversity (π) and recombination rate (ρ) across different subpopulations. **b** Comparisons of the average number of dTEs in the selective windows (i.e., ranked in top 5% of 100 kb F_{ST} windows for SNPs) and in windows randomly selected from each permutation for 500 independent permutations between subpopulations. **c** Number of dTEs showing signatures of selection (henceforth termed “selected dTEs”) between *Or* and *Osi*, *Or* and *Osj*, *Osi* and *Osj*. F_{ST} outlier based on TE variations were indicated as loci showing signatures of selection. **d** Fold enrichment of selected dTEs in each TE family by the Fisher’s exact test. The purple symbols refer to significant enrichment or depletion and gray symbols indicate no significant enrichment. Unknown refers to LTR/unknown family. **e** Positions of functional genes harboring selected dTEs in their genic regions on chromosomes 1–6.

Chromosomes 7–12 see **Fig. S3g**. Genic regions included the gene body, regions of 2kb upstream (henceforth termed “promoter”) and 2kb downstream (henceforth termed “downstream”) of the gene body. Functional genes harboring selected dTEs in rice domestication and differentiation processes are displayed in purple and green font, respectively. The shared genes harboring selected dTEs between domestication and differentiation processes are displayed in blue font. Differences in the gene expression between subpopulations were tested using the Student’s *t*-test. *** $P < 0.001$, ** $P < 0.01$, and * $P < 0.05$. **f** A 150 bp dTE (*Tourist* MITE) insertion occurred at 262 bp upstream of *LIP19* in *Osj* accessions. **g-h** Differences in the expression level of *LIP19* (**g**) and cold tolerance (**h**) between *Osi* accessions with the ancestral state ($n=131$ and $n=122$, respectively) and *Osj* accessions with the derived state ($n=58$ and $n=54$, respectively) of the 150 bp dTE INS. Significance was tested by the Student’s *t*-test, *** $P < 0.001$. **i** Phenotypes of NH242 (an *Osj* accession) and NH181 (an *Osi* accession) under control (**left**) and cold stress treatments (**right**). For the cold stress treatment, 10-day-old seedlings were transferred to 4°C for 72h and recovered for 4 days in a greenhouse at 30°C. **j** Differences in the absolute value of latitude of the accessions with this dTE INS (derived state) and those without this dTE INS (ancestral state) near *LIP19*. Significance was tested by the Wilcoxon-Mann-Whitney test, ** $P < 0.01$. **k** Phylogenetic tree based on the sequence alignments of the promoter and gene body regions of *LIP19* across all accessions in the present study. Accessions in different subpopulations are indicated by different colors. The pink and green dots indicate in accessions with (derived state) and without (ancestral state) the dTE, respectively..




Fig. 4 TE variations associated with gene expression and agricultural traits.

a The distribution of linkage disequilibrium values (LD, R^2) between dTEs and SNPs/InDels within 50kb of the dTEs. For each dTE, the maximum R^2 with adjacent SNPs/InDels (within 50kb on either side) was recorded. The dashed line indicates $R^2=0.70$. **b** Percentage of dTEs overlapping with genic regions. Genic regions include gene body (CDS, 3'UTR, 5'UTR, and intron), and the regions within 2 kb upstream (henceforth termed "promoter") and 2kb downstream (henceforth termed "downstream") of the gene body. Unknown refers to LTR/unknown family. "All" refers to the percentage of dTEs in the genic regions of all dTE loci identified in the present study, and "NIP" refers the percentage of TEs overlapping with genic regions in the Nipponbare genome. **c** Number of eGenes associated with dTE and SNP variants. eGenes are genes whose

expression is significantly associated with dTE and SNP variants. **d** Manhattan plot of *OsRbohB* expression level and the dTE variants and SNPs (**top**). The leading dTE (328 bp) insertion (INS) associated with *OsRbohB* expression level is indicated (**bottom**). **e-f** Differences in the expression level of *OsRbohB* (**e**) and the thousand grain weight (**f**) between accessions with the ancestral state of the dTE INS event (n=125 and n=121, respectively) and those with the derived state (both n=61). Significance was tested by the Student's *t*-test, ****P* < 0.001. **g** Transcriptional activation assays by cotransfecting rice protoplasts with the constructs shown in the upper panel. Error bars represent the mean ± SD of three biological replicates. **h** GWAS for seed setting rate under cold stress using the TE and SNP datasets for the *Os* accessions. A locus on chromosome 6, which was identified by TEs but not by SNPs, was significantly associated with seed setting rate under cold stress. The green triangles represent SNPs and pink dots represent dTEs. The most strongly associated dTE was a 1.0 kb dTE (*Gypsy*) deletion (DEL) event in the promoter of *LOC_Os06g28970*. **i-j** Comparison of seed setting rate under cold stress (**i**) and the expression level of *LOC_Os06g28970* (**j**) between accessions with (i.e., derived state, n=16 and n=32, respectively) and without (i.e., ancestral state, n= 83 and n=169, respectively) the dTE DEL event. Significance was tested by the Student's *t*-test, ***P* < 0.01, **P* < 0.05.